

## SHORT-COMMUNICATION

## A comparison of horizontal versus vertical camera placement to detect feral cats and mustelids

Margaret Nichols<sup>1\*</sup>, Alistair S. Glen<sup>2</sup>, Patrick Garvey<sup>3</sup> and James Ross<sup>1</sup><sup>1</sup>Centre for Wildlife Management and Conservation, Lincoln University, Canterbury, New Zealand<sup>2</sup>Landcare Research, Private Bag 92170, Auckland 1142, New Zealand<sup>3</sup>School of Biological Science, Tamaki campus, University of Auckland, Private Bag 92019, Auckland 1142, New Zealand

\*Author for correspondence (Email: Maggie.Nichols@lincolnuni.ac.nz)

Published online: 30 September 2016

**Abstract:** Invasive predators are a threat to biodiversity in New Zealand. However, they are often difficult to monitor because of the animals' cryptic, mobile behaviour and low densities. Camera traps are increasingly being used to monitor wildlife, but until recently have been used mainly for large species. We aimed to determine the optimal camera alignment (horizontal or vertical) for detecting feral cats (*Felis catus*) and mustelids (*Mustela furo*, *M. erminea* and *M. nivalis*). We deployed 20 pairs of cameras, each pair with one horizontal and one vertical camera. We compared the number of photos of target species, non-target species, and false triggers (i.e. camera triggered with no animal present) between camera orientations. Horizontally oriented cameras captured approximately 1.5 times as many images of the target species compared with vertically oriented cameras, and also detected more non-target animals. Orientation did not have a significant effect on the number of false triggers.

**Keywords:** camera trapping; feral ferret; invasive species; *Mustela* spp.; stoat

## Introduction

Invasive mammalian predators are among the greatest threats to New Zealand's biodiversity (Krull et al. 2015), but can be difficult to monitor due to their highly cryptic nature, and in some cases (i.e. feral cats) low densities (Glen et al. 2013). In recent decades, various methods have been used to assess mammal abundance and distribution, including trapping, hair snags, spotlight counts, scat surveys, camera traps and tracking tunnels (Gompper et al. 2006; Long et al. 2007a, b; Pickerell et al. 2014; Lazenby et al. 2015). In New Zealand, tracking tunnels have been the most commonly used non-lethal method for monitoring small mammals such as rodents and mustelids (King & Edgar 1977; Brown et al. 1996). Although there are indeed many successful monitoring methods available for small to medium-sized mammals, over the last 20 years attention has turned towards camera traps as an effective research tool (Rowcliffe & Carbone 2008). Since camera traps are remotely triggered and impervious to most weather conditions, they may be left for long periods of time for monitoring purposes (Long et al. 2007a; Meek et al. 2014a). Camera traps may also have higher detection rates than some other monitoring techniques such as tracking tunnels and live capture traps (Sam 2011), and have the potential to identify uniquely marked individuals (Heilbrun et al. 2003; Long et al. 2007a, b; Sam 2011).

Numerous studies have used camera traps for large mammals such as leopard (*Uncia uncia*), jaguar (*Panthera onca*) and tiger (*Panthera tigris*) (Karanth et al. 2004; Jackson et al. 2006; Kelly et al. 2008; Tobler et al. 2008; Wang & Macdonald 2009), but only a handful have examined the optimal specifications for small to medium-sized species (e.g. De Bondi et al. 2010; Glen et al. 2013; Bischof et al. 2014). There is a wide range of variables associated with camera traps, from trigger settings to sensor types as well as data

analysis methods (De Bondi et al. 2010; Meek et al. 2014a). Additionally, camera orientation, along with height from the ground, detection zone, distance from a lure (if used), and the size of the target species must all be considered when deploying camera traps (Smith & Coulson 2012; Glen et al. 2013; Taylor et al. 2013; Meek et al. 2014a). Camera traps are usually oriented horizontally at a height to accommodate the size of the target species (Smith & Coulson 2012). De Bondi et al. (2010) tested an alternative approach by placing cameras vertically, angled at 90 degrees facing towards the ground to capture photos from above the target – a technique now known as vertical orientation (Smith & Coulson 2012). This method has the advantage of standardising the size of the camera's detection zone, but orientation may also affect the success of certain camera traps in detecting animals that encounter the traps.

Smith and Coulson (2012) compared vertical and horizontal orientation for two Australian marsupials, potoroos (*Potorous tridactylus*, 660–1640 g; Norton et al. 2011) and bandicoots (*Isodon obesulus*, >1 kg; De Milliano et al. 2016). They found that vertically oriented cameras had a detection probability for these target species up to five times greater than horizontal cameras. Taylor et al. (2013) performed a similar study with bandicoots, potoroos and pademelons (*Thylogale stigmatica*, 4–7 kg; Macqueen et al. 2009). However, this study found horizontally oriented cameras had detection probabilities 2.5 times greater than vertically oriented cameras. These studies varied in both deployment and set-up methods.

We aimed to compare the effectiveness of horizontal and vertical camera trap orientations for detecting feral cats (*Felis catus*) and mustelids (feral ferrets *Mustela furo*, stoats *M. erminea* and weasels *M. nivalis*). Like the marsupials mentioned above, these species range in size, with typical cats weighing 1–5 kg, ferrets 600–1200 g, stoats 200–325 g

and weasels 55–125 g (King 2005). We compared the number of photos of target species between cameras with these two orientations, along with the number of false triggers (when cameras were triggered without capturing an image of an animal), the total number of photos taken throughout the study (including target species, non-target species and false triggers), and the number of independent encounters (Brook et al. 2012) with individuals of the target species (as distinct from repeated images of the same animal).

## Methods

### Study area and field methods

The study was conducted on Toronui Station, a pastoral property in Hawke's Bay, North Island, New Zealand (39°0' S, 176°46' E). Toronui Station (1600 ha) is mainly covered by introduced pasture grass, with fragments of native beech forest (*Fuscospora solandri*), on both high country and lowland paddocks (300–1000 m above sea level). Fence lines were often hedged with pines (*Pinus radiata*) as windbreaks for livestock, which included red deer (*Cervus elaphus*), sheep (*Ovis aries*) and cattle (*Bos taurus*).

From 20 January to 24 March 2014, 20 pairs of cameras were placed along existing monitoring transects. Paired camera trap sites were spaced 2.4 km apart on average, with a minimum separation of 700 m. We placed cameras at the ecotones of forest fragments wherever possible, to increase predator detection

rates (Meek et al. 2014a). Two cameras were placed 1.5 m apart at each station. One camera was placed on a steel fence post facing vertically towards the ground from a height of 1.5 m. The other was set horizontally, 7 cm from the ground (as measured to the base of the camera) and attached to a tree or wooden stake (Fig. 1). As part of a concurrent trial, European rabbit meat (*Oryctolagus cuniculus*) and ferret odour (towel bedding from a male ferret's enclosure) lures were separately contained in two perforated vials, and set directly beneath the vertical-facing camera. This design allowed the lure vials to be within the field of view of both cameras.

We primarily used Reconyx Hyperfire PC900 trail cameras (Reconyx Inc., Holmen, Wisconsin, USA), but also LTL Acorn 5210A (Shenzen LTL Acorn Electronics Co., Ltd, Shenzhen, Guangdong, China), M990i (Moultrie, Calera, Alabama, USA) and Bushnell (Bushnell Outdoor Products, Overland Park, Kansas, USA) (see Table 1 for camera types, specifications and settings). All cameras were chosen for their infrared flash, which is likely to be less conspicuous to cats than a white flash (Glen et al. 2013; but see also Meek et al. 2014b). Vegetation was cleared to a height of 5 cm where necessary to provide a clear view of animals in the detection zone and to avoid possible false triggers caused by moving branches or foliage (Kelly & Holub 2008; Taylor et al. 2013). Cameras were checked after 4 weeks, and batteries, memory cards (4–8 GB) and scent lures were replaced. Photos were uploaded onto an external hard drive according to their site number and orientation. All photographed animals were recorded in an Excel™ file along with any false triggers, following the methods of Allen (2014).



**Figure 1.** Setup of horizontal and vertical cameras, Toronui Station, New Zealand, in 2014. Reconyx cameras are shown below; camera models and settings are given in Table 1.

**Table 1.** Camera specifications and settings used at Toronui Station, New Zealand, in 2014.

	Camera type			
	Reconyx ®	LTL Acorn ®	Moultrie ®	Bushnell ®
Trigger speed (seconds)	0.2	0.8	0.69	0.2
Recovery time (seconds)	0.5	1	5	1
Flash range (metres)	15	15	15	24
Sensor	PIR	PIR	PIR	PIR
Light source	Infrared flash	Infrared flash	Infrared flash	Infrared flash
Sense level (normal, high, low)	Normal	Normal	Normal	Normal
Number of photos per trigger	3	3	3	3
Number of cameras	24	10	4	2

### Data analysis

Photographs were classed as either: (1) target species, (2) non-target species, or (3) false trigger events. To increase the sample size, we pooled cats and mustelids for analysis simply as ‘target species’. We plotted histograms of the elapsed time between successive photographs of the target species to isolate encounters with an individual animal from repeated observations of the same individual (Brook et al. 2012). The average time between consecutive photographs of cats was <10 minutes, indicating these to be repeat detections. Therefore, we assumed photographs taken >30 minutes apart were ‘independent encounters’ representing separate individuals, except for individuals that could be identified reliably (e.g. by coat pattern). Similarly, on the basis of the activity patterns of mustelids, (consecutive photographs <5 minutes apart) we assumed encounters >15 minutes apart were independent.

We used the software program GENSTAT version 15 (VSN International 2011) to create generalised linear mixed-effects models. A Poisson error distribution was used as we had continuous count data. To assess the performance of the two camera orientations at capturing target species, camera orientation (vertical or horizontal) was fitted as a fixed effect, and camera type and the camera monitoring stations were random effects. We used likelihood ratio tests to compare models with each of four response variables (numbers of target species photos, independent encounters with target species, all photos and false triggers) to the corresponding null model without an orientation parameter.

### Results

Data from 36 of the original 40 cameras were used. One camera was damaged by livestock, one was damaged by flooding and two cameras had memory cards filled to capacity, due to false triggers and livestock. The cameras detected 79 independent encounters with cats (50 on the horizontal cameras and 29 on vertical cameras), 45 independent encounters with

stoats (25 horizontal and 20 vertical), and two independent encounters with ferrets (horizontal only). There were also 23 independent encounters with target species that were detected by both camera orientations. No weasels were detected. Non-target species (83% of all photos taken) included house mouse (*Mus musculus*), ship rat (*Rattus rattus*), Norway rat (*R. norvegicus*), brushtail possum (*Trichosurus vulpecula*), European hedgehog (*Erinaceus europaeus occidentalis*), European rabbit, feral pig (*Sus scrofa*), Eurasian blackbird (*Turdus merula*), house sparrow (*Passer domesticus*) and silvereye (*Zosterops lateralis*).


Horizontally oriented cameras yielded significantly more photos of target species compared with vertical cameras ( $\chi^2 = 4.54$ , d.f. = 1, 15,  $P = 0.05$ ) (Table 2, Fig. 2a). Horizontally placed cameras also captured significantly more independent encounters with target species than did the vertical cameras ( $\chi^2 = 5.55$ , d.f. = 1, 15.4,  $P = 0.032$ ) (Fig. 2b), and significantly more photos in total (false triggers, target and non-target species) ( $\chi^2 = 15.67$ , d.f. = 1, 22.1,  $P = 0.001$ ) (Table 2, Fig. 2c). However, orientation did not significantly affect the number of false triggers ( $\chi^2 = 0.41$ , d.f. = 1, 16.7,  $P = 0.53$ ) (Table 2, Fig. 2d). Vertical cameras often provided clearer images than horizontal cameras of the coat patterns of cats. However, the large body size of cats relative to the camera’s field of view meant that 63% of cats photographed by vertical cameras were partially outside the frame. The corresponding proportion for horizontal cameras was 36%.

### Discussion

Our results showed that horizontally placed cameras were more effective at detecting the target species, i.e. cats and mustelids combined. Smith and Coulson (2012) found that the wider field of view associated with the horizontal cameras decreased detection rates for small to medium-sized species. There were differences in camera set-up including distance

**Table 2.** Number of photos of target and non-target species, and false trigger events obtained from cameras with horizontal and vertical alignment, at Toronui Station, New Zealand, in 2014.

Orientation	Target species (stoats, ferrets, cats)	Non-target species	False triggers	Total photos
Horizontal	832	22 117	3746	26 695
Vertical	571	11 478	2013	14 062



**Figure 2.** (a) Number of photos of target species, (b) independent encounters with a target species, (c) total photos over all, and (d) false triggers by cameras with horizontal and vertical alignment. Camera models and settings are given in Table 1.

from the horizontal camera to the lure (3 m, 2 m, and 1.5 m respectively). There was also a difference in camera settings (i.e. continuous triggering (Smith & Coulson 2012) vs bursts of three images with a forced delay (Taylor et al. 2013).

There has been some debate over the optimal camera trap orientation for identifying individuals of a species (Smith & Coulson 2012; Taylor et al. 2013). De Bondi et al. (2010) observed that vertical cameras assisted in the species identification of mammals (smaller than cats) that fitted entirely in a camera’s field of view. In contrast, although we found that cats’ coat patterns were clearest in photographs taken directly beneath vertical cameras, full coat identifications would potentially have been difficult because the cats were

often only partially in the fields of view of the vertical cameras.

In conclusion, our study shows that horizontal cameras are likely to detect more cats and mustelids than vertical cameras mounted at 1.5 m. However, should a study’s aim be to identify individuals through coat patterns, further investigation into the utility of vertical camera orientation may be necessary. While vertical cameras may help identify animals, cameras must have a sufficiently wide field of view to capture complete images of the target species. Future studies could test vertical cameras raised >1.5 m from the ground to broaden the field of view and compensate for the larger size of feral cats, to improve coat identification.

## Acknowledgements

This study was funded by the Hawke's Bay Regional Council and by core funding to Landcare Research from the Ministry of Business, Innovation and Employment. Many thanks to R. Dickson and S. Cave for their technical support and to D. Shaw for site access to Toronui Station. We also greatly appreciate the helpful comments from Deb Wilson and two anonymous reviewers on a previous version of the manuscript.

## References

- Allen B 2014. How to collect, store and query pest animal data: a tutorial for practitioners. In: Gentle M ed. Program and abstracts 16th Australasian Vertebrate Pest Conference, Brisbane. Pp. 87.
- Bischof R, Ali H, Kabir M, Hameed S, Nawaz M 2014. Being the underdog: an elusive small carnivore uses space with prey and time without enemies. *Journal of Zoology* 219: 40–48.
- Brook LA, Johnson CN, Ritchie EG 2012. Effects of predator control on behaviour of an apex predator and indirect consequences for mesopredator suppression. *Journal of Applied Ecology* 49: 1278–1286.
- Brown K, Moller H, Innes J, Alterio N 1996. Calibration of tunnel tracking rates to estimate relative abundance of ship rats (*Rattus rattus*) and mice (*Mus musculus*) in a New Zealand forest. *New Zealand Journal of Ecology* 20: 271–275.
- DeBondi N, White JG, Stevens M, Cooke R 2010. A comparison of the effectiveness of camera trapping and live trapping for sampling terrestrial small-mammal communities. *Wildlife Research* 37: 456–465.
- De Milliano J, Di Stefano J, Courtney P, Temple-Smith P, Coulson G 2016. Soft-release versus hard-release for reintroduction of an endangered species: an experimental comparison using eastern barred bandicoots (*Perameles gunnii*). *Wildlife Research* 43: 1–12.
- Glen AS, Cockburn S, Nichols M, Ekanayake J, Warburton B 2013. Optimising camera traps for monitoring small mammals. *PloS one* 8(6): e67940.
- Gompper ME, Kays RW, Ray JC, Lapoint SD, Bogan DA, Cryan JR 2006. A comparison of noninvasive techniques to survey carnivore communities in northeastern North America. *Wildlife Society Bulletin* 34: 1142–1151.
- Heilbrun RD, Silvy NJ, Tewes ME, Peterson MJ 2003. Using automatically triggered cameras to individually identify bobcats. *Wildlife Society Bulletin*: 748–755.
- Jackson RM, Roe JD, Wangchuk R, Hunter DO 2006. Estimating snow leopard population abundance using photography and capture-recapture techniques. *Wildlife Society Bulletin* 34: 772–781.
- Karanth KU, Chundawat RS, Nichols JD, Kumar NS 2004. Estimation of tiger densities in the tropical dry forests of Panna, Central India, using photographic capture-recapture sampling. *Animal Conservation* 7: 285–290.
- Kelly MJ, Holub EL 2008. Camera trapping of carnivores: trap success among camera types and across species, and habitat selection by species, on Salt Pond Mountain, Giles County, Virginia. *Northeastern Naturalist* 15: 249–262.
- Kelly MJ, Noss AJ, Di Bitetti MS, Maffei L, Arispe RL, Paviolo A, De Angelo CD, Di Blanco YE 2008. Estimating puma densities from camera trapping across three study sites: Bolivia, Argentina, and Belize. *Journal of Mammalogy* 89: 408–418.
- King CM 2005. The handbook of New Zealand mammals. 2nd edn. Melbourne, Oxford University Press. 610 p.
- King CM, Edgar R 1977. Techniques for trapping and tracking stoats (*Mustela erminea*); a review, and a new system. *New Zealand Journal of Zoology* 4: 193–212.
- Krull C, Galbraith JA, Glen AS, Nathan HW 2015. Invasive vertebrates in Australia and New Zealand. In: Stow A, MacLean N, Holwell GI eds. *Austral ark: the state of wildlife in Australia and New Zealand*. Cambridge, Cambridge University Press. Pp. 197–226.
- Lazenby BT, Mooney NJ, Dickman CR 2015. Effects of low-level culling of feral cats in open populations: a case study from the forests of southern Tasmania. *Wildlife Research* 41: 407–420.
- Long RA, Donovan TM, Mackay P, Zielinski WJ, Buzas JS 2007a. Comparing scat detection dogs, cameras, and hair snares for surveying carnivores. *Journal of Wildlife Management* 71: 2018–2025.
- Long RA, Donovan TM, Mackay P, Zielinski WJ, Buzas JS 2007b. Effectiveness of scat detection dogs for detecting forest carnivores. *Journal of Wildlife Management* 71: 2007–2017.
- Macqueen P, Goldizen AW, Seddon JM 2009. Response of a southern temperate marsupial, the Tasmanian pademelon (*Thylogale billardierii*), to historical and contemporary forest fragmentation. *Molecular ecology* 18: 3291–3306.
- Meek P, Ballard G, Claridge A, Kays R, Moseby K, O'Brien T, O'Connell A, Sanderson J, Swann D, Tobler M 2014a. Recommended guiding principles for reporting on camera trapping research. *Biodiversity and Conservation* 23: 2321–2343.
- Meek PD, Ballard G-A, Fleming PJ, Schaefer M, Williams W, Falzon G 2014b. Camera traps can be heard and seen by animals. *PloS one* 9: e110832.
- Norton MA, Claridge AW, French K, Prentice A 2011. Population biology of the long-nosed potoroo (*Potorous tridactylus*) in the Southern Highlands of New South Wales. *Australian Journal of Zoology* 58: 362–368.
- Pickerell GA, O'Donnell CF, Wilson DJ, Seddon PJ 2014. How can we detect introduced mammalian predators in non-forest habitats? A comparison of techniques. *New Zealand Journal of Ecology* 38: 86–102.
- Rowcliffe JM, Carbone C 2008. Surveys using camera traps: are we looking to a brighter future? *Animal Conservation* 11: 185–186.
- Sam S 2011. New monitoring and control tools for simultaneously managing possums, rats and mice in New Zealand. Unpublished PhD thesis, Lincoln, Lincoln University. 149 p.
- Smith JK, Coulson G 2012. A comparison of vertical and horizontal camera trap orientations for detection of potoroos and bandicoots. *Australian Mammalogy* 34: 196–201.
- Taylor BD, Goldingay RL, Lindsay JM 2013. Horizontal or vertical? Camera trap orientations and recording modes for detecting potoroos, bandicoots and pademelons. *Australian Mammalogy* 36: 60–66.
- Tobler MW, Carrillo-Percastegui SE, Leite Pitman R, Mares R, Powell G 2008. An evaluation of camera traps for inventorying large- and medium-sized terrestrial rainforest mammals. *Animal Conservation* 11: 169–178.
- VSN International 2011. *GenStat for Windows*. 14th edn.

Hemel Hempstead, UK, VSN International.

Wang SW, Macdonald DW 2009. The use of camera traps for estimating tiger and leopard populations in the high altitude mountains of Bhutan. *Biological Conservation* 142: 606–613.

Editorial board member: Deb Wilson

Received 20 April 2016; accepted 15 August 2016